

Target™ Detachable Coils family product offering



All deliverable through Excelsior™ SL-10™ and XT-17™ Microcatheters



360 shape



3D shape




Helical shape

Secondary OD (mm)	360 shape						3D shape			Helical shape		
	Target XL Standard	Target XL Soft	Standard	Soft	Ultra	Nano	Standard	Target XL Soft	Ultra	Nano		
24	50											
22	50											
20	50	50										
18	50	50										
16	50	50										
15			40									
14	50	50	30	30								
13			30	30								
12	45	45	30	30								
11			30	30								
10	40	40	30	30			30	40				
9	30	30	30, 20	30, 20				30				
8	30	30	30, 20	30, 20			25	30				
7	20	20	30, 20, 15	30, 20, 15			15	20				
6	20	20	30, 20, 15	30, 20, 15, 10			15	20				
5		15, 10	20, 15	20, 15, 10	15, 10		10	15				
4.5				12	10				8			
4		12, 8	15, 10, 8	15, 10, 8, 6	15, 10, 8, 6		8	15, 8, 6				
3.5				10	8			6				
3		9, 6	8, 6	10, 8, 6	10, 8, 6, 4	6	6	10, 8, 6, 4	8, 6, 4			
2.5					4	4		6, 4, 3	6, 4, 3			
2		6, 3		4	6, 4, 3	4, 3		8, 6, 4, 3, 2, 1	6, 4, 3, 2			
1.5						4, 3, 2			4, 3, 2, 1			
1						3, 2			3, 2, 1			


Size and softness

Target XL™ Coils




Target XL Standard

Target XL Soft




Standard

2x softer




Soft

2x softer



Ultra

2x softer



Nano™

10-Size Coils

Target™ Detachable Coil

See package insert for complete indications, contraindications, warnings and instructions for use.

INTENDED USE / INDICATIONS FOR USE

Target Detachable Coils are intended to endovascularly obstruct or occlude blood flow in vascular abnormalities of the neurovascular and peripheral vessels.

Target Detachable Coils are indicated for endovascular embolization of:

- Intracranial aneurysms
- Other neurovascular abnormalities such as arteriovenous malformations and arteriovenous fistulae
- Arterial and venous embolizations in the peripheral vasculature

THIS DOCUMENT IS INTENDED SOLELY FOR THE USE OF HEALTHCARE PROFESSIONALS.

A physician must always rely on his or her own professional clinical judgment when deciding whether to use a particular product when treating a particular patient. Stryker does not dispense medical advice and recommends that physicians be trained in the use of any particular product before using it in a procedure. The information presented is intended to demonstrate the breadth of Stryker product offerings. A physician must always refer to the package insert, product label and/or instructions for use before using any Stryker product. Products may not be available in all markets because product availability is subject to the regulatory and/or medical practices in individual markets. Please contact your Stryker representative if you have questions about the availability of Stryker products in your area. The Stryker products listed above are CE marked according to the Medical Device Directive 93/42/EEC.